

Walgreens

-PE Product Labels –

CONVENIENT RECLOSING TAB

Well at Walgreens
WALGREENS PHARMACIST RECOMMENDED¹

DAYTIME Wal-Flu[®]
Severe Cold & Cough

Drug Facts

Active Ingredients (in each packet)

Acetaminophen 650 mg	Pain reliever/Fever reducer
Dextromethorphan hydrobromide 20 mg	Cough suppressant
Phenylephrine hydrochloride 10 mg	Nasal decongestant

Uses

- Temporarily relieves:
 - minor aches and pains
 - headache
 - minor sore throat pain
 - nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
- Temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 packets in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen or 3 or more alcoholic drinks every day while using this product.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription and nonprescription). Ask a doctor or pharmacist before using with other drugs if you are not sure.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema, asthma, or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product

- do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- symptoms do not get better or worsen
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- take every 4 hours; not to exceed 6 packets in 24 hours or as directed by a doctor.
- adults and children 12 years of age and over: dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- children under 12 years of age: consult a doctor.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 8 mg, sodium 20 mg
- phenylephrine HCl: contains phenylephrine 14 mg per packet
- store at controlled room temperature 20-25°C (68-77°F). Protect from excessive heat and moisture.

Inactive ingredients: Acetaminophen, aspartame, citric acid, FD&C Blue No. 1, FD&C red No. 40, maltodextrin, menthol, natural flavors, sodium citrate, starch and sugar.

DO NOT USE IF SEALED PACKET IS TORN OR BROKEN

Well at Walgreens
WALGREENS PHARMACIST RECOMMENDED¹

NEW DAYTIME Wal-Flu[®]
Severe Cold & Cough

**Acetaminophen / Pain Reliever / Fever Reducer
Dextromethorphan HBr / Cough Suppressant
Phenylephrine HCl / Nasal Decongestant**

Relieves:

- Nasal & Sinus Congestion
- Cough • Body Ache
- Sore Throat Pain
- Headache • Fever



BERRY
Infused with Menthol & Green Tea Flavors

Compare to Theraflu[®] Daytime Severe Cold & Cough active ingredients¹¹

6 PACKETS **SEE NEW WARNINGS INFORMATION**

¹¹Walgreens Pharmacist Survey Study, November 2012.
¹²This product is not affiliated with Novartis Consumer Health, Inc., owner of the registered trademark Theraflu[®] Daytime Severe Cold & Cough.

Questions or comments? 1-800-925-4733

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

walgreens.com ©2013 Walgreen Co.
MADE IN CANADA

ITEM 279236



3 11917 15938 6

C-3110-30508-6W
ORG0413-F

[illegible]

ORG code is an internal Periscope code, that will be updated by Periscope prior

[illegible][illegible]




1-800-828-6478




Compare to the active ingredients
in Therma Multi-Symptom
Severe Cold*

MULTI-SYMPTOM

Severe Cold

**ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER /
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT /
PHENYLEPHRINE HCl / NASAL DECONGESTANT**

Dye Free

Aspartame Free **Sodium Free**

- Relieves nasal congestion,
sore throat pain, cough,
headache, body ache & fever



Honey Lemon
Infused with Citrus Flavors
& with No Real Fruit

6
PACKETS

*NOT DELIVERED BY WALGREEN CO.
 206 WALWAT RD., DEERFIELD, IL 60015
 100% SATISFACTION GUARANTEED
 walgreens.com
 ©2021 Walgreen Co.

81281 870381 053002C-00005-0



3 11917 20156 8

SA 705942121C

Lot:

Exp:

READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE.
KEEP CARTON FOR REFERENCE. DO NOT DISCARD.

Drug Facts

Active Ingredients (in each packet)

Acetaminophen 500 mg. Pain reliever/fever reducer
Diphenhydramine HCl 25 mg. Antihistamine/cough suppressant
Phenylephrine HCl 10 mg. Nasal decongestant

Purposes

Uses

- Temporarily relieves these symptoms due to a cold:
 - runny nose
 - nasal congestion
 - cough
 - throat irritation
 - fever
 - aches and pains
 - headache
 - sinus and sinus congestion

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours.

Do not take with other products containing acetaminophen.

Do not use if you are allergic to any of the ingredients.

Always read the label on all medicines you take. Symptoms may include:

- skin rash, hives, or itching
- swelling, especially of the face, lips, or tongue

If a skin reaction occurs, stop use and seek medical help right away.

Some throat warning: If your throat is sore, use for no more than 2 days. If accompanied by

difficulty swallowing, fever, or breathing, stop use and seek medical help right away.

Do not use if you are under 12 years of age. If you are allergic to acetaminophen,

do not use. If you are allergic to diphenhydramine, do not use. If you are

not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

If you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- heart disease
- heart failure
- high blood pressure
- thyroid disease
- diabetes
- glaucoma

prostate problems due to an enlarged prostate gland

breathing problems such as emphysema or chronic bronchitis

cough that occurs after a cold (postnasal drip)

cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are:

- taking medicines or medical tests
- taking the blood thinning drug, warfarin

When using this product:

- do not exceed recommended dosage
- avoid alcohol drinks
- marked drowsiness may occur

Do not drink alcohol or use grapefruit juice while taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Stop use and ask a doctor if:

- newborns, infants, or preschoolers
- newborns or infants
- newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

newborns or infants

Drug Facts (continued)

- Do not use if your cold or flu symptoms get worse or last more than 7 days.
- Cough comes back or occurs with rash or wheezing that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Prompt medical attention is critical for adults as well as for children.

Even if you do not notice any signs or symptoms.

Do not use if you are allergic to any of the ingredients.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a

doctor or pharmacist before taking this product.

Do not use if you are pregnant or breastfeeding.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

Walgreens



Walgreens

Compare to the active ingredients
in Therflu Nighttime Severe
Cold & Cough¹



NCC 0883-91048

NIGHTTIME

Severe Cold & Cough

COLD & COUGH

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DIPHENHYDRAMINE HCl / ANTIHISTAMINE /
COUGH SUPPRESSANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

Aspartame Free Sodium Free

- Relieves nasal congestion, cough, body ache, sore throat pain, runny nose, sneezing, headache & fever



Honey Lemon
Infused with chamomile & white tea flavors

6

PACKETS



W30R08122-F

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2021 Walgreen Co.

ITEM 978380 W00000-0000-0



54100WALGP-C

Lot:

Exp:

Drug Facts

Active ingredients (in each packet)

Acetaminophen 650 mg	Pain reliever/Fever reducer
Diphenhydramine hydrochloride 25 mg	Antihistamine/Cough suppressant
Phenylephrine hydrochloride 10 mg	Nasal decongestant

Uses

- Temporarily relieves:
 - minor aches and pains
 - headache
 - minor sore throat pain
 - nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
 - runny nose
 - sneezing
 - itchy, watery eyes due to hay fever
 - itchy nose and throat
- Temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 packets in 24 hours, which is the maximum daily amount, with other drugs containing acetaminophen, 3 or more alcoholic drinks every day while using this product. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription and nonprescription). Ask a doctor or pharmacist before using with other drugs if you are not sure.
- with any other product containing diphenhydramine, even one used on the skin.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- glaucoma
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema, asthma, or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema.

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- symptoms do not get better or worsen
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health care professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- take every 4 hours; not to exceed 6 packets in 24 hours or as directed by a doctor.
- adults and children 12 years of age and over: dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- children under 12 years of age: consult a doctor.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 7 mg, sodium 46 mg
- phenylephrine: contains phenylalanine 13 mg per packet
- store at controlled room temperature 20-25°C (68-77°F). Protect from excessive heat and moisture.

Inactive ingredients: Acesulfame K, aspartame, caramel color, citric acid, D&C Yellow 10, maltodextrin, natural flavors, sodium citrate, starch and sugar.

Questions or comments?
1-800-925-4733

CONVENIENT RECLOSING TAB

NIGHTTIME
Wal-Flu®
Severe Cold & Cough

WALGREENS PHARMACIST RECOMMENDED[†]

Well at Walgreens

NIGHTTIME
Wal-Flu®
Severe Cold & Cough

Acetaminophen / Pain Reliever / Fever Reducer
Diphenhydramine HCl / Antihistamine / Cough Suppressant
Phenylephrine HCl / Nasal Decongestant

Relieves:

- Nasal congestion • Cough
- Body ache • Sore throat pain
- Runny nose • Sneezing
- Headache • Fever

HONEY LEMON FLAVOR
Infused with Chamomile & White Tea Flavors

Compare to Theraflu® Nighttime Severe Cold & Cough active ingredients^{††}

6 PACKETS **SEE NEW WARNINGS INFORMATION**

[†]Walgreens Pharmacist Survey Study, November 2012
^{††}This product is not affiliated with Novartis Consumer Health, Inc., owner of the registered trademark Theraflu®.

DO NOT USE IF SEALED PACKETS ARE TORN OR BROKEN

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

Walgreens
100% SATISFACTION GUARANTEED

walgreens.com © 2013 Walgreen Co.
MADE IN CANADA

ITEM 499403

3 11917 10758 5

C-3109-30508-6W-2 REV0813-RF

Well at Walgreens

WALGREENS PHARMACIST RECOMMENDED*

COMBO PACK

NON-DROWSY • DAYTIME

Sinus Relief

**Acetaminophen / Sinus Headache & Pain
Phenylephrine HCl / Sinus Pressure & Congestion**

32
Liquid Caps

COMBO PACK

NIGHTTIME

Sinus Relief

**Acetaminophen / Sinus Headache & Pain
Doxylamine Succinate / Runny Nose
Phenylephrine HCl / Sinus Pressure & Congestion**

16
Liquid Caps

Compare to Vicks® DayQuil™ & NyQuil™ Sinex® Sinus Relief LiquidCaps® active ingredients**

48 TOTAL LIQUID CAPS

B9845

JFBG687J

Nighttime Severe Cold & Cough

Drug Facts

Active ingredients (in each packet) **Purpose**
Acetaminophen 650 mg **Pain Reliever / Fever Reducer**
Diphenhydramine hydrochloride 25 mg **Antihistamine / Cough Suppressant**
Phenylephrine hydrochloride 10 mg **Nasal Decongestant**

Uses

- Temporarily relieves:
 - minor aches and pains • headache • minor sore throat pain
 - nasal and sinus congestion • cough due to minor throat and bronchial irritation • runny nose • sneezing • itchy, watery eyes due to hay fever • itchy nose and throat
- Temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 8 packets in 24 hours, which is the maximum daily amount. Do not take more than 8 packets in 24 hours, which is the maximum daily amount. Do not take more than 8 packets in 24 hours, which is the maximum daily amount. Do not take more than 8 packets in 24 hours, which is the maximum daily amount.

Do not use

- with any other drug containing acetaminophen (prescription and nonprescription)
- Ask a doctor or pharmacist before using with other drugs if you are not sure
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease • heart disease • high blood pressure • thyroid disease
- diabetes • blood clotting due to an enlarged prostate gland
- a bleeding problem such as emphysema, asthma, or chronic bronchitis
- cough that occurs with too much phlegm (mucus) • cough that lasts or is chronic such as occurs with smoking, asthma or emphysema.

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers • taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks • marked drowsiness may occur • alcohol, sedatives and tranquilizers may increase drowsiness • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children.

Stop use and ask a doctor if

- innuminate, diarrhea, or sleeplessness occurs • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur
- symptoms do not get better or worse • pain, cough or nasal congestion gets worse or lasts more than 7 days • cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

Daytime Severe Cold & Cough

Drug Facts

Active ingredients (in each packet) **Purpose**
Acetaminophen 650 mg **Pain Reliever / Fever Reducer**
Dextromethorphan hydrobromide 20 mg **Cough Suppressant**
Phenylephrine hydrochloride 10 mg **Nasal Decongestant**

Uses

- Temporarily relieves:
 - minor aches and pains • minor sore throat pain
 - headache • nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
- Temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 8 packets in 24 hours, which is the maximum daily amount. Do not take more than 8 packets in 24 hours, which is the maximum daily amount. Do not take more than 8 packets in 24 hours, which is the maximum daily amount. Do not take more than 8 packets in 24 hours, which is the maximum daily amount.

Do not use

- with any other drug containing acetaminophen (prescription and nonprescription)
- Ask a doctor or pharmacist before using with other drugs if you are not sure
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease • heart disease • high blood pressure • thyroid disease
- diabetes • blood clotting due to an enlarged prostate gland
- a bleeding problem such as emphysema, asthma, or chronic bronchitis
- cough that occurs with too much phlegm (mucus) • cough that lasts or is chronic such as occurs with smoking, asthma or emphysema.

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers • taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks • marked drowsiness may occur • alcohol, sedatives and tranquilizers may increase drowsiness • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children.

Stop use and ask a doctor if

- innuminate, diarrhea, or sleeplessness occurs • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur
- symptoms do not get better or worse • pain, cough or nasal congestion gets worse or lasts more than 7 days • cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

Daytime

Drug Facts

Active ingredients (in each packet) **Purpose**
Acetaminophen 650 mg **Pain Reliever / Fever Reducer**
Dextromethorphan hydrobromide 20 mg **Cough Suppressant**
Phenylephrine hydrochloride 10 mg **Nasal Decongestant**

Uses

- Temporarily relieves:
 - minor aches and pains • minor sore throat pain
 - headache • nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
- Temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 8 packets in 24 hours, which is the maximum daily amount. Do not take more than 8 packets in 24 hours, which is the maximum daily amount. Do not take more than 8 packets in 24 hours, which is the maximum daily amount. Do not take more than 8 packets in 24 hours, which is the maximum daily amount.

Do not use

- with any other drug containing acetaminophen (prescription and nonprescription)
- Ask a doctor or pharmacist before using with other drugs if you are not sure
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease • heart disease • high blood pressure • thyroid disease
- diabetes • blood clotting due to an enlarged prostate gland
- a bleeding problem such as emphysema, asthma, or chronic bronchitis
- cough that occurs with too much phlegm (mucus) • cough that lasts or is chronic such as occurs with smoking, asthma or emphysema.

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers • taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks • marked drowsiness may occur • alcohol, sedatives and tranquilizers may increase drowsiness • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children.

Stop use and ask a doctor if

- innuminate, diarrhea, or sleeplessness occurs • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur
- symptoms do not get better or worse • pain, cough or nasal congestion gets worse or lasts more than 7 days • cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

COMBO PACK

NEW

Wal-Flu®

Severe Cold & Cough

Acetaminophen / Pain Reliever / Fever Reducer
Dextromethorphan HBr / Cough Suppressant
Phenylephrine HCl / Nasal Decongestant

Relieves:

- Nasal & sinus congestion
- Cough • Body ache
- Sore throat pain
- Headache • Fever

6 PACKETS

12 TOTAL PACKETS

SEE NEW WARNINGS INFORMATION

Lot code area - keep text/graphics free

Questions or comments? 1-800-925-4733

DISTRIBUTED BY: WALGREEN CO.
200 WALMART RD., DEERFIELD, IL 60015
ITEM 367905
100% SATISFACTION GUARANTEED
walgreens.com ©2014 Walgreen Co.
MADE IN CANADA

COMBO PACK

NIGHTTIME

Wal-Flu®

Severe Cold & Cough

Acetaminophen / Pain Reliever / Fever Reducer
Diphenhydramine HCl / Antihistamine / Cough Suppressant
Phenylephrine HCl / Nasal Decongestant

Relieves:

- Nasal congestion • Cough
- Runny nose • Sneezing
- Body ache • Sore throat pain
- Headache • Fever

6 PACKETS

12 TOTAL PACKETS

SEE NEW WARNINGS INFORMATION

Lot code area - keep text/graphics free

Questions or comments? 1-800-925-4733

DISTRIBUTED BY: WALGREEN CO.
200 WALMART RD., DEERFIELD, IL 60015
ITEM 367905
100% SATISFACTION GUARANTEED
walgreens.com ©2014 Walgreen Co.
MADE IN CANADA

Drug Facts

Active ingredients (in each packet)

Acetaminophen 650 mg	Pain reliever/Fever reducer
Diphenhydramine hydrochloride 25 mg	Antihistamine/Cough suppressant
Phenylephrine hydrochloride 10 mg	Nasal decongestant

Uses

- Temporarily relieves:
 - minor aches and pains
 - headache
 - minor sore throat pain
 - nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
 - runny nose
 - sneezing
 - itchy, watery eyes due to hay fever
 - itchy nose and throat
- Temporarily reduces fever

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 packets in 24 hours, which is the maximum daily amount, or with other drugs containing acetaminophen or 3 or more alcoholic drinks every day while using this product. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- in a child under 4 years of age
- with any other drug containing acetaminophen (prescription and nonprescription). Ask a doctor or pharmacist before using with other drugs. If you are not sure
- with any other product containing diphenhydramine, even one used on the skin
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- glaucoma
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema, asthma, or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema.

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- symptoms do not get better or worsen
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health care professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- take every 4 hours; not to exceed 6 packets in 24 hours or as directed by a doctor.
- adults and children 12 years of age and over: dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- children under 12 years of age: consult a doctor.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 10 mg, sodium 19 mg
- phenylketonurics: contains phenylalanine 11 mg per packet
- store at controlled room temperature 20-25°C (68-77°F). Protect from excessive heat and moisture.

Inactive ingredients: Acesulfame K, aspartame, calcium phosphate dibasic, citric acid, D&C Yellow 10, maltodextrin, natural flavor, silicon dioxide, sodium citrate dihydrate.

DO NOT USE IF SEALED PACKET IS TORN OR BROKEN.

Questions or comments?
1-800-925-4733

CONVENIENT RECLOSING TAB

SUGAR FREE

NIGHTTIME

Wal-Flu®

Severe Cold & Cough

WALGREENS PHARMACIST RECOMMENDED¹

NEW

Well at Walgreens

SUGAR FREE

NIGHTTIME

Wal-Flu®

Severe Cold & Cough

Acetaminophen / Pain Reliever / Fever Reducer
Diphenhydramine HCl / Antihistamine / Cough Suppressant
Phenylephrine HCl / Nasal Decongestant

Temporarily relieves:

- Nasal congestion
- Cough
- Body ache
- Sore throat pain
- Runny nose
- Sneezing
- Headache
- Fever

NATURAL HONEY LEMON FLAVOR

Compare to Theraflu® Sugar Free Nighttime Severe Cold & Cough^{††}

6 PACKETS

¹Walgreens Pharmacist Survey Study, November 2012.
^{††}This product is not affiliated with Novartis Consumer Health, Inc., owner of the registered trademark Theraflu®.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

ITEM 376865

100% SATISFACTION GUARANTEED

walgreens.com ©2014 Walgreen Co.
MADE IN CANADA

3 11917 17294 1

C-3111-30508-6W

Walgreens
Way to Well
Commitment

Support local preventive wellness with Walgreens. With each purchase of Walgreens Brand Health & Wellness Product, 1¢ goes to the Walgreens Way to Well Fund[®] up to \$3 million annually. Visit Walgreens.com/waytowell

READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE. KEEP CARTON FOR REFERENCE. DO NOT DISCARD

Drug Facts

Active Ingredients (in each packet)

Active Ingredients (in each packet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Pheniramine maleate 20 mg	Antihistamine
Phenylephrine hydrochloride 10 mg	Nasal Decongestant

Uses

- temporarily relieves these symptoms due to a cold:
 - minor aches and pains • headache • minor sore throat/pain • nasal congestion
- temporarily reduces fever
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose • sneezing • itchy nose and throat • itchy, watery eyes

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take • more than 6 packets in 24 hours, which is the maximum daily amount; • with other drugs containing acetaminophen; • 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin redness • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a physician promptly.

Do not use • in a child under 4 years of age • if you are allergic to acetaminophen • with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease • heart disease • high blood pressure • thyroid disease • diabetes
- glaucoma • trouble urinating due to an enlarged prostate gland • cough that occurs with too much phlegm (mucus) • cough that lasts or is chronic such as occurs with smoking, asthma or emphysema • a sodium-restricted diet.

Ask a doctor or pharmacist before use if you are taking

- sedatives or tranquilizers • the blood thinning drug warfarin

When using this product • do not exceed recommended dosage • avoid alcoholic drinks • may cause drowsiness • alcohol, sedatives, and tranquilizers may increase drowsiness • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs • pain or nasal congestion gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days. (continued)

Drug Facts (continued)

- redness or swelling is present • new symptoms occur • These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- take every 4 hours; not to exceed 6 packets in 24 hours or as directed by a doctor.

Age	Dose
children under 4 years of age	do not use
children 4 to under 12 years of age	do not use unless directed by a doctor
adults and children 12 years of age and over	one packet

• dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.

• If using a microwave, add contents of one packet to 8 oz. of cool water; stir briefly before and after heating. Do not overheat.

Other Information

- each packet contains: sodium 66 mg
- store at controlled room temperature 20-25°C (68-77°F). Protect from heat and moisture.

Inactive ingredients

Cholic acid, DMC Yellow 13, Ethylcellulose, FD&C Red 40, Flavor, Microcrystalline cellulose, Povidone, Silicon dioxide, Sodium citrate, Starch, Sugar and Vitamin C.

Questions or comments? 1-800-925-4733

DO NOT USE IF SEALED PACKETS ARE TORN OR BROKEN

¹Walgreens Pharmacist Survey Study, November 2014.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2015 Walgreen Co.
MADE IN CANADA

CONVENIENT RECLOSING TAB

Well at Walgreens
WALGREENS PHARMACIST RECOMMENDED¹

Wal-Flu®
Cold & Sore Throat


Well at Walgreens
WALGREENS PHARMACIST RECOMMENDED¹

Wal-Flu®
Cold & Sore Throat

Acetaminophen / Pain Reliever / Fever Reducer
Pheniramine Maleate / Antihistamine
Phenylephrine HCl / Nasal Decongestant

Relieves:

- Nasal congestion
- Body ache
- Sore throat pain
- Runny nose
- Headache
- Fever

6 PACKETS  **LEMON FLAVOR**

EV Code


Lot/Exp

C-3173-30508-6W-3

REV1115-RF5

Source: *Journal of the American Statistical Association*, 92(439), 1031-1042.

NDC 0363-5430-09



W30RG0821-F2
REV1122

Do not print area
Lot and
Exp Date Only

ITEM 603348 W00000-0000-0
3 11917 06812 1

B-2201-485-08-HR
REV1220A48508

Drug Facts (continued)
mfcrcystalline cellulose, polydextrose, polyethylene glycol, silicon dioxide, stearic acid, titanium dioxide, triacetin
1-800-426-9391
(Questions or comments?)

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. Johnson & Johnson Corporation, owner of the registered trademark Benadryl® Allergy Plus Congestion.

Drug Facts (continued)
When using this product
do not exceed recommended dosage
excitability may occur, especially in children
marked drowsiness may occur
alcohol, sedatives, and tranquilizers may increase drowsiness
avoid alcoholic beverages
use caution when driving a motor vehicle or operating machinery
Stop use and ask a doctor if
symptoms do not improve within 7 days or are accompanied by fever
nervousness, dizziness, or sleeplessness occur
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions ■ do not take more than directed
adults and children 12 years and over
take 1 tablet every 4 hours
do not take more than 6 tablets in 24 hours
children under 12 years: do not use
Other information
each tablet contains: calcium 25 mg
IS OPENED OR BLISTER IS TORN OR BROKEN
TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE
store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
see end flap for expiration date and lot number
Inactive ingredients
croscarmellose sodium, aluminum lake, hypromellose, magnesium stearate, disbasic calcium phosphate dihydrate, FD&C blue #1

Drug Facts
Active ingredients
(in each tablet)
Diphenhydramine HCl 25 mgAntihistamine
Phenylephrine HCl 10 mgNasal decongestant
Purpose
COMPLETE PRODUCT INFORMATION
KEEP OUTER PACKAGE FOR
Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
runny nose ■ sneezing ■ nasal congestion
itching of the nose or throat ■ itchy, watery eyes
temporarily relieves these symptoms due to the common cold:
runny nose ■ sneezing ■ nasal congestion
temporarily relieves sinus congestion and pressure
Do not use
if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug.
If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
with any other product containing diphenhydramine, even one used on skin
Ask a doctor before use if you have
difficulty in urination due to enlargement of the prostate gland ■ diabetes ■ thyroid disease
heart disease ■ glaucoma ■ high blood pressure
a breathing problem such as emphysema or chronic bronchitis
sedatives or tranquilizers.
Ask a doctor or pharmacist before use if you are taking

how2recycle.info
PAPER BOX
TRAY
TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING
DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2021 Walgreen Co.
50844 REV1220A48508

Walgreens Allergy Relief


Walgreens
Compare to the active ingredients in Benadryl® Allergy Plus Congestion*

Allergy Relief PLUS CONGESTION

DIPHENHYDRAMINE HCl 25 mg / ANTIHISTAMINE
PHENYLEPHRINE HCl 10 mg / NASAL DECONGESTANT

Congestion Relief
• Relief of sinus congestion & pressure, runny nose, sneezing, itchy throat & itchy, watery eyes

24



No Print/No Varnish
Lot and Expiration No





Compare to Children's Mucinex® Multi-Symptom Cold active ingredients†

children's
MULTI-SYMPTOM
Cold

DEXTROMETHORPHAN HBr /
COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCl /
NASAL DECONGESTANT


ALCOHOL FREE


- Relief from: stuffy nose, cough & chest congestion
- Breaks up mucus

AGES
4-12
YEARS



4 FL OZ (118 mL) **BERRY FLAVOR**







Compare to Children's Mucinex® Multi-Symptom Cold active ingredients†


children's
MULTI-SYMPTOM
Cold

DEXTROMETHORPHAN HBr /
COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCl /
NASAL DECONGESTANT


ALCOHOL FREE


- Relief from: stuffy nose, cough & chest congestion
- Breaks up mucus

AGES
4-12
YEARS



4 FL OZ (118 mL) **BERRY FLAVOR**





children's
MULTI-SYMPTOM
Cold

DEXTROMETHORPHAN HBr /
COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCl /
NASAL DECONGESTANT

Drug Facts

Active Ingredients (in each 5 mL)

Dextromethorphan HBr 5 mg.....Cough suppressant

Guaifenesin 100 mg.....Expectorant

Phenylephrine HCl 2.5 mg.....Nasal decongestant

Purposes

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the intensity of coughing
- the impulse to cough to help your child get to sleep
- nasal congestion due to a cold
- stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product do not use more than directed

Stop use and ask a doctor if

- your child gets nervous, dizzy or sleepless
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter

Drug Facts (continued)

Age	Dose
children 6 years to under 12 years	10 mL every 4 hours
children 4 years to under 6 years	5 mL every 4 hours
children under 4 years	do not use

Other Information

- each 5 mL contains: sodium 3 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not refrigerate
- dosing cup provided

Inactive Ingredients anhydrous citric acid, edetate disodium, FD&C red #40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?
1-800-719-9260

Gluten Free

Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey
†This product is not manufactured or distributed by Reckitt Benckiser, the distributor of Children's Mucinex® Multi-Symptom Cold.

DOSAGE CUP PROVIDED


PARENTS:
Learn about teen medicine abuse.
www.StopMedicineAbuse.org

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

walgreens.com ©2017 Walgreen Co.

ITEM 374549 W10106-1217-L



3 11917 17279 8

LOT NO.

EXP.

• 83926 94 C4



Compare to Children's Dimetapp®
Cold & Cough active ingredients††

NDC 0363-0708-26

children's
COUGH & COLD
Wal-Tap®
DM

BROMPHENIRAMINE MALEATE /
ANTIHISTAMINE
DEXTROMETHORPHAN HBr /
COUGH SUPPRESSANT
PHENYLEPHRINE HCl /
NASAL DECONGESTANT

ALCOHOL FREE

- Relieves nasal congestion;
runny nose; itchy, watery eyes;
sneezing & coughing

6 YEARS & OLDER

4 FL OZ (118 mL) GRAPE FLAVOR





Compare to Children's Dimetapp®
Cold & Cough active ingredients††

NDC 0363-0708-26

children's
COUGH & COLD
Wal-Tap®
DM

BROMPHENIRAMINE MALEATE /
ANTIHISTAMINE
DEXTROMETHORPHAN HBr /
COUGH SUPPRESSANT
PHENYLEPHRINE HCl /
NASAL DECONGESTANT

ALCOHOL FREE

- Relieves nasal congestion;
runny nose; itchy, watery eyes;
sneezing & coughing

6 YEARS & OLDER

4 FL OZ (118 mL) GRAPE FLAVOR

ORG0717-F

Walgreens
children's
Wal-Tap®
COUGH & COLD

DO NOT USE IF PRINTED NECK BAND IS
BROKEN OR MISSING

MA 5.1 L DECONGESTANT
PHE 10 mg / 5 mL
DM 5 mg / 5 mL
PH 5 mg / 5 mL
MA 5.1 L DECONGESTANT

Drug Facts

Active Ingredients (in each 10 mL)

Brompheniramine maleate, USP 2 mg.....Antihistamine
Dextromethorphan HBr, USP 10 mg.....Cough suppressant
Phenylephrine HCl, USP 5 mg.....Nasal decongestant

Purposes

- temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- temporarily restores freer breathing through the nose

Uses

- temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- do not use more than directed
- may cause marked drowsiness
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Other Information

- each 10 mL contains: sodium 4 mg
- store at 20°-25°C (68°-77°F)

Inactive Ingredients: anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution

Questions or comments?

1-800-719-9260

Gluten Free

Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey

†† This product is not manufactured or distributed by Pfizer, distributor of Children's Dimetapp® Cold & Cough.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

DOSAGE CUP PROVIDED

ITEM 126858 W10106-1217-L

3 11917 10249 8

LOT NO.

EXP.

96726 94 C8

NDC 0363-0666-40

Walgreens

Compare to the active ingredients in Vicks® DayQuil®††

DAYTIME • NON-DROWSY Cold & Flu

ACETAMINOPHEN /
ACHES / FEVER / SORE THROAT
DEXTROMETHORPHAN HBr / COUGH
PHENYLEPHRINE HCl /
NASAL CONGESTION

Multi-Symptom

- Pain reliever, fever reducer, cough suppressant & nasal decongestant
- Antihistamine free
- Alcohol free

12 FL OZ (355 mL)
: 65640 94 F7

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.

ITEM 163066 W10107-0522-F

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

Gluten Free
W30RG0222-F

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses ■ temporarily relieves common cold/flu symptoms:
■ cough due to minor throat and bronchial irritation
■ nasal congestion ■ minor aches and pains
■ sore throat ■ headache ■ fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

PEEL BACK AT CORNER FOR MORE INFORMATION

11917 14623 2

: 65640 94 B1

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ if you have ever had an allergic reaction to this product or any of its ingredients

ADHESIVE AREA
• NO VARNISH • NO TYPE

Drug Facts (continued)


Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed - see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor

ADHESIVE AREA
• NO VARNISH • NO TYPE





Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Maximum Strength Mucinex® FAST-MAX® Cold & Flu. DISTRIBUTED BY: WALGREEN CO. DEERFIELD, IL 60015 100% SATISFACTION GUARANTEED walgreens.com ©2023 Walgreen Co.

PARENTS: Learn about your medicine visit www.dapdrmedusa.org

PEEL BACK TAB TO READ COMPLETE DRUG FACTS AND INFORMATION

ITEM 875753 W00000-0000-0
B-042-45 ORG
50844 ORG022304245
3 11917 03987 5
W3ORG0923-F

Drug Facts

Active ingredients (in each 20 mL)

Active ingredients (in each 20 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - stuffy nose
 - minor aches and pains
 - headache
 - cough
 - sore throat
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

No Print / No Varnish Area
Lot # and Exp. Info

Drug Facts (continued)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- diabetes
- cough that occurs with too much phlegm (mucus)
- liver disease
- thyroid disease
- heart disease
- difficulty in urination due to enlargement of the prostate gland
- high blood pressure
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Drug Facts (continued)

Stop use and ask a doctor if

- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- do not take more than 6 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- adults and children 12 years and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years: do not use

Other information

- each 20 mL contains: sodium 10 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous citric acid, disodium edetate, FD&C blue #1, FD&C red #40, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose, xanthan gum

Questions or comments? Call 1-800-426-9391

3147

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- heart disease
- diabetes
- thyroid disease
- liver disease
- high blood pressure

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

¹Our pharmacists recommend the Walgreens brand.

We invite you to compare to national brands.

²This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® DayQuil® Cold & Flu.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2022 Walgreen Co.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

ITEM 852225 W00000-0000-0



W3ORG0922-F
50844 REV0323A01202 REV0423

No Print / No Varnish Area
Lot # and Exp. Info

Do Not Take
Daytime and
Nighttime
Products at the
Same Time.



Walgreens

Compare to the active ingredients
in Vicks® DayQuil® Cold & Flu¹



DAY

Cold & Flu

ACETAMINOPHEN /
PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr /
COUGH SUPPRESSANT
PHENYLEPHRINE HCl /
NASAL DECONGESTANT

- Relieves headaches, fever,
sore throat, minor aches
& pains, cough
- Nasal congestion

Ages 6 Years and Over

12 FL OZ (355 mL)



Menthol
flavor

NDC 0363-9912-02

Drug Facts

TAMPER EVIDENT: DO NOT USE IF PRINTED
NECK WRAP IS BROKEN OR MISSING

**Active ingredients
(in each 15 mL)**

	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
- sore throat
- nasal congestion
- minor aches and pains
- cough due to minor throat and bronchial irritation
- headache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
 - child takes more than 5 doses in 24 hours
 - taken with other drugs containing acetaminophen
 - adult has 3 or more alcoholic drinks every day while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if the user has

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- do not exceed 4 doses per 24 hours
- mL = milliliter
- only use the dose cup provided

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children under 6 years	do not use

Other information

- each 15 mL contains: sodium 13 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients anhydrous citric acid, FD&C yellow #6, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sodium saccharin, sorbitol, sucralose

Questions or comments? 1-800-426-9391

DAY & NIGHT PACK[illegible]



Compare the active ingredients in Vicks[®] VapoGel[®] Cold & Flu & Vicks[®] Hyalinel[®] Cold & Flu



COMPARE THE ACTIVE INGREDIENTS

DAY

Cold & Flu

ACETAMINOPHEN /
PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr /
COUGH SUPPRESSANT
PHENYLEPHRINE HCl /
NASAL DECONGESTANT

- Relieves headaches, fever, sore throat, minor aches & pains, cough
- Nasal congestion

• Ages 6 Years and Over



Menthol
flavor

NIGHT

Cold & Flu

ACETAMINOPHEN /
PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr /
COUGH SUPPRESSANT
Doxylamine Succinate /
ANTIHISTAMINE

- Relieves headaches, fever, sore throat, minor aches & pains, sneezing, cough
- Runny nose

• Ages 12 Years and Over



Cherry
flavor

2 - 12 FL OZ (355 mL) BOTTLES / TOTAL 24 FL OZ (710 mL)

[illegible]

PARENTS:
Read this information carefully.

Your pharmacist(s) recommend the Wegovy[®] brand, because you have to compare to national brands. This product is one manufacturer or distributed by The Procter & Gamble Company, owner of the registered trademark Wegovy[®], Qsulfam[®] C&F, and Flus.

This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Wegovy[®], Qsulfam[®] C&F, and Flus.



ITEM# 42578 W00020-0000
 1171712592

W-2501-012013-02TW
 50864 REV0235A01201302
 W30G00262-F REV0523

No Print / No Varnish Area
Lot # and Exp. Info

3150

48 TOTAL CAPLETS

Drug Facts
Active ingredients (in each caplet)
 Acetaminophen 325 mg... Pain reliever/fever reducer
 Dextromethorphan HBr 10 mg... Cough suppressant
 Guaifenesin 200 mg... Expectorant
 Phenylephrine HCl 5 mg... Nasal decongestant

Drug Facts (continued)
Uses
 ■ temporarily relieves common cold and flu symptoms:
 ■ minor aches and pains ■ sinus congestion and pressure ■ nasal congestion
 ■ sore throat ■ fever ■ headache
 ■ cough due to minor throat and bronchial irritation

Drug Facts (continued)
 ■ temporarily restores free breathing through the nose ■ promotes nasal and/or sinus drainage
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive ■ reduces swelling of nasal passages

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ blisters ■ rash ■ skin reddening
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
 ■ liver disease ■ thyroid disease ■ diabetes
 ■ heart disease ■ high blood pressure
 ■ difficulty in urination due to enlargement of the prostate gland

Drug Facts (continued)
 ■ cough that occurs with too much phlegm (mucus)
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
When using this product do not exceed recommended dosage.

Stop use and ask a doctor if
 ■ nervousness, dizziness, or sleeplessness occur
 ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ new symptoms occur
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed
 ■ do not take more than 8 caplets in 24 hours
 ■ adults and children 12 years and over: take 2 caplets with water every 4 hours
 ■ children under 12 years: ask a doctor

Other information ■ each caplet contains: sodium 3 mg ■ **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN** ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number

Inactive ingredients corn starch, croscopollose, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Drug Facts (continued)
Questions or comments?
 1-800-426-9391

DISTRIBUTED BY: WALGREEN CO.
 200 WILMOT RD., DEERFIELD, IL 60015
 100% SATISFACTION GUARANTEED
 ©2021 Walgreen Co.

ITEM 329615 W00000-0000-0
 3 19171 6389 5

Do not remove this caplet from the blister pack until you are ready to use it. This product is not manufactured or distributed by the Project & Sample Pharmacy. Walgreen's Severe Cold & Flu.
 50844 REV0722A64008

W3ORG1121-F
 REV1122

No Print/No Varnish
 Lot # & Exp. Area

TAMPER EVIDENT:
 DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

PARENTS:
 Learn about teen medicine abuse
 www.StopMedicineAbuse.org



B-2201-640-08-HRR
 REV0722A64008



NDC 0383-0640-08

Compare to the active ingredients in Vicks® DayQuil® Severe Cold & Flu®

DAY Severe Cold & Flu

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER
 DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
 GUAIFENESIN 200 mg / EXPECTORANT
 PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

Maximum Strength

• Relieves headache, fever, sore throat, minor aches & pains, chest congestion, thins & loosens mucus, nasal congestion, sinus pressure & cough

24 CAPLETS



ACTUAL SIZE



Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. *This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Maximum Strength Mucinex® NIGHTSHIFT® Severe Cold & Flu. DISTRIBUTED BY: WALGREEN CO., DEERFIELD, IL 60015 100% SATISFACTION GUARANTEED walgreens.com ©2023 Walgreen Co.

PARENTS: 60644 ORG102307845 B-078-45 ORG

FEEL BACK TAB TO READ COMPLETE DRUG FACTS AND INFORMATION

W30RG0923-F
ITEM 872975 W00000-0000-0

11917 03714

Drug Facts

Active ingredients (in each 20 mL) Purpose

Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Phenylephrine HCl 10 mg.....	Nasal decongestant
Tripolidine HCl 2.5 mg.....	Antihistamine

Uses ■ temporarily relieves these common cold and flu symptoms: ■ nasal congestion ■ sore throat ■ minor aches and pains ■ cough ■ sinus congestion and pressure ■ sneezing ■ itching of the nose or throat ■ runny nose ■ itchy, watery eyes due to hay fever ■ headache ■ temporarily reduces fever ■ controls cough to help you get to sleep

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ rash ■ blisters ■ skin reddening. If a skin reaction occurs, stop use and

No Print / No Varnish Area Lot # and Exp. Info

Drug Facts (continued)

seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ if you have ever had an allergic reaction to this product or any of its ingredients ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ diabetes ■ cough that occurs with too much phlegm (mucus) ■ heart disease ■ liver disease ■ difficulty in urination due to enlargement of the prostate gland ■ high blood pressure ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ glaucoma ■ a breathing problem such as emphysema or chronic bronchitis ■ thyroid disease

Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

When using this product ■ do not exceed recommended dosage ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ avoid alcoholic beverages

Drug Facts (continued)

■ use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ new symptoms occur ■ nervousness, dizziness, or sleeplessness occur ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed ■ do not take more than 4 doses in any 24-hour period ■ mL = milliliter ■ only use the dose cup provided ■ adults and children 12 years and over: 20 mL in dosing cup provided every 4 hours ■ children under 12 years: do not use

Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

Inactive ingredients anhydrous citric acid, FD&C blue #1, FD&C red #40, FD&C yellow #6, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions or comments? 1-800-426-9391

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg... Pain reliever/fever reducer
Dextromethorphan HBr 10 mg... Cough suppressant
Doxylamine succinate 6.25 mg... Antihistamine
Phenylephrine HCl 5 mg... Nasal decongestant

Purpose

■ temporarily relieves common cold and flu symptoms:
■ sinus congestion and pressure
■ sore throat ■ fever ■ headache
■ nasal congestion ■ minor aches and pains
■ cough due to minor throat and bronchial irritation ■ runny nose and sneezing

Drug Facts (continued)

Uses

■ temporarily relieves common cold and flu symptoms:
■ sinus congestion and pressure
■ sore throat ■ fever ■ headache
■ nasal congestion ■ minor aches and pains
■ cough due to minor throat and bronchial irritation ■ runny nose and sneezing

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ blisters ■ rash ■ skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

■ liver disease ■ diabetes ■ thyroid disease
■ heart disease ■ glaucoma
■ difficulty in urination due to enlargement of the prostate gland ■ high blood pressure
■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

Inactive ingredients black iron oxide, corn starch, croscopolvidone, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Do not print Lot and Exp Only

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ blisters ■ rash ■ skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

■ liver disease ■ diabetes ■ thyroid disease
■ heart disease ■ glaucoma
■ difficulty in urination due to enlargement of the prostate gland ■ high blood pressure
■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

Inactive ingredients black iron oxide, corn starch, croscopolvidone, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Do not print Lot and Exp Only

Walgreens

Compare to Vicks® NyQuil® Severe Cold & Flu active ingredients††

NDC 0363-6770-08

NIGHTTIME

Severe Cold & Flu

**ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
DOXYLAMINE SUCCINATE / ANTIHISTAMINE
PHENYLEPHRINE HCl / NASAL DECONGESTANT**

MAXIMUM STRENGTH

• Relieves headache, fever, sore throat, minor aches & pains, nasal/sinus congestion & sinus pressure, sneezing, runny nose & cough

24
CAPLETS

ACTUAL SIZE

24
CAPLETS

ACTUAL SIZE

50844 REV0519B67708

DISTRIBUTED BY: WALGREEN CO.,
200 WILMOT RD., DEERFIELD, IL 60015

ITEM 907683 W00000-0000-0

11 917 17797

7

TAMPER EVIDENT: DO NOT USE IF
PACKAGING IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING

Walgreens Prescription Drug
††This product is not manufactured or
distributed by The Pacer Company
Company, owner of the registered trademark
Vicks® NyQuil® Severe Cold & Flu.

ORG0219-F

PHARMACIST

Walgreens

Trusted since 1901™

RECOMMENDED

Health expertise
you rely on.™

8-2201-677-08-HR
REV0519B67708

Questions or comments?
1-800-426-9391

Walgreens

Compare to Mucinex® Fast-Max® Cold, Flu & Sore Throat active ingredients††

MUCUS RELIEF

NDC 0363-1209-30

Cold, Flu & Sore Throat

ACETAMINOPHEN /
PAIN RELIEVER & FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

MAXIMUM STRENGTH

- Relieves headache, fever, sore throat, nasal & chest congestion
- Thins & loosens mucus
- Controls cough
- 12 years & older

6 FL OZ (180 mL)

: 16930 94 F1

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

DO NOT USE IF PRINTED NECKLAMP IS CRACKED OR MISSING

Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey
DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2017 Walgreen Co.
ORG0118-F



Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

Uses ■ temporarily relieves these common cold and flu symptoms:
■ cough ■ nasal congestion ■ minor aches and pains ■ headache
■ sore throat ■ temporarily reduces fever ■ temporarily promotes
nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin
bronchial secretions to rid the bronchial passageways of bothersome
mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver
damage may occur if you take ■ more than 4,000 mg of acetaminophen
in 24 hours ■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause
severe skin reactions. Symptoms may
include: ■ skin reddening ■ blisters
■ rash. If a skin reaction occurs, stop
use and seek medical help right away.

**PEEL CORNER TO READ COMPLETE
DRUG FACTS AND INFORMATION**

: 16930 94 B1

Drug Facts (continued)

If pregnant or breast-feeding, ask a health
professional before use.
Keep out of reach of children.

Overdose warning: In case of overdose, get
medical help or contact a Poison Control Center
right away (1-800-222-1222). Quick medical
attention is critical for adults as well as for children
even if you do not notice any signs or symptoms.

Directions ■ do not take more than
directed (see Overdose warning) ■ do not
take more than 5 doses in any 24-hour period
■ measure only with dosing cup provided
■ do not use dosing cup with other products
■ dose as follows or as directed by a doctor
■ adults and children 12 years of age and older:
20 mL in dosing cup provided every 4 hours
■ children under 12 years of age: do not use

Other Information

■ each 20 mL contains: sodium 7 mg
■ dosing cup provided ■ store at 20-25°C
(68-77°F) ■ do not refrigerate

Inactive ingredients anhydrous citric acid,
benzyl alcohol, edetate disodium, FD&C blue #1,
FD&C red #40, flavor, glycerin, propylene glycol,
purified water, sodium benzoate, sodium citrate,
sorbitol solution, sucralose, triethyl citrate,
xanthan gum

Questions or comments?
1-800-719-9260

†† This product is not manufactured
or distributed by Reckitt.
Bencik, distributor of
Mucinex® Fast-Max® Cold,
Flu & Sore Throat.

Drug Facts (continued)

Sore throat warning: If sore throat is severe,
persists for more than 2 days, is accompanied or
followed by fever, headache, rash, nausea, or
vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing
acetaminophen (prescription or nonprescription).
If you are not sure whether a drug contains
acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine
oxidase inhibitor (MAOI) (certain drugs for depression,
psychiatric, or emotional conditions, or Parkinson's
disease), or for 2 weeks after stopping the MAOI drug.
If you do not know if your prescription drug contains
an MAOI, ask a doctor or pharmacist before taking
this product. ■ if you have ever had an allergic
reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ liver
disease ■ heart disease ■ diabetes ■ high
blood pressure ■ thyroid disease ■ trouble
urinating due to an enlarged prostate gland
■ persistent or chronic cough such as occurs with
smoking, asthma, chronic bronchitis, or emphysema
■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are
taking the blood thinning drug warfarin

**When using this product do not use more than
directed**

Stop use and ask a doctor if ■ nervousness,
dizziness or sleeplessness occur ■ pain, nasal
congestion, or cough gets worse or
lasts more than 7 days ■ fever gets
worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur ■ cough
comes back or occurs with rash or
headache that lasts. These could be
signs of a serious condition.

NDC 0363-8005-45

Walgreens

Compare to the active ingredients
in Maximum Strength Mucinex®
FAST-MAX® Cold, Flu & Sore Throat[†]

**Cold, Flu
& Sore Throat**

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

Maximum Strength

- Relieves body pain, headache, fever & sore throat
- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus
- Ages 12 years & over

6 FL OZ (177 mL)

F-005-45 ORG

PARENTS:
Learn about our products and more at
www.DrugFactsOnline.org

DISTRIBUTED BY:
WALGREEN CO.
DEERFIELD, IL 60015
100% SATISFACTION
GUARANTEED
walgreens.com
©2023 Walgreen Co.

50844 ORG052300545
B-005-45 ORG
ITEM 769447 W00000-0000-0

3 11917 03411 9
W3ORG0923-F

[†]Our pharmacists recommend the Walgreens brand.
We invite you to compare to national brands.
^{††}This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Maximum Strength Mucinex® FAST-MAX® Cold, Flu & Sore Throat.

Drug Facts

Active ingredients (in each 20 mL)

Active ingredients (in each 20 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses ■ temporarily relieves these common cold and flu symptoms:

- minor aches and pains
- sinus congestion and pressure
- stuffy nose ■ nasal congestion
- sore throat ■ cough ■ headache
- temporarily promotes nasal and/or sinus drainage
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**No Print / No Varnish Area
Lot # and Exp. Info**

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- rash ■ blisters ■ skin reddening. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- liver disease ■ heart disease ■ diabetes
- difficulty in urination due to enlargement of the prostate gland ■ high blood pressure ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ thyroid disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Drug Facts (continued)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ new symptoms occur
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed

- do not take more than 6 doses in any 24-hour period
- mL = milliliter ■ only use the dose cup provided
- dose as follows or as directed by a doctor
- adults and children 12 years and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years: do not use

Other information ■ each 20 mL contains: sodium 10 mg ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

Inactive ingredients anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose

Questions or comments? 1-800-426-9391



DAYTIME • NON-DROWSY

Cold & Flu



NIGHTTIME

Cold & Flu

DAY & NIGHT PACK

WDC 0903-0567-02



Compare to the active ingredients in
Vicks® DayQuil® & Vicks® NyQuil®[†]

DAYTIME
NON-DROWSY

**Cold
& Flu**

ACETAMINOPHEN / ACHES /
FEVER / SORE THROAT
DEXTROMETHORPHAN HBr /
COUGH
PHENYLEPHRINE HCl /
NASAL CONGESTION

Multi-Symptom

- Pain reliever, fever reducer,
cough suppressant
& nasal decongestant
- Antihistamine free
- Alcohol free



NIGHTTIME

**Cold
& Flu**

ACETAMINOPHEN / ACHES /
FEVER / SORE THROAT
DEXTROMETHORPHAN HBr /
COUGH
DOXYLAMINE SUCCINATE /
SNEEZING / RUNNY NOSE

Multi-Symptom

- Pain reliever, fever reducer,
cough suppressant
& antihistamine
- ALCOHOL 10%

2 - 12 FL OZ (355 mL) BOTTLES / TOTAL 24 FL OZ (1.5 pt) (710 mL)

DO NOT USE IF PRINTED NECKBAND IS
BROKEN OR MISSING

Gluten Free

Empty & Replace Cap



PAPER
BOX

PLASTIC
BOTTLE

PLASTIC
CUP

how2recycle.info

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

[†]Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
^{**}These products are not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks® NyQuil® and DayQuil®.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015


100% SATISFACTION
GUARANTEED

ITEM 165449 W10108-0722-L



Nighttime Cold & Flu

Drug Facts

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

Uses temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning
 - only use the dose cup provided
 - do not exceed 4 doses per 24 hrs
- | | |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 6 hrs |
| children 4 to under 12 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- each 30 mL contains: sodium 38 mg
- store at 20-25°C (68-77°F)

Inactive ingredients alcohol, anhydrous citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments? 1-800-719-9260

Daytime Cold & Flu

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning
 - only use the dose cup provided
 - do not exceed 4 doses per 24 hrs
- | | |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs | 15 mL every 4 hrs |
| children 4 to under 6 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- each 15 mL contains: sodium 7 mg
- store at 20-25°C (68-77°F)

Inactive ingredients butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments? 1-800-719-9260

W3ORG0422-F




DAYTIME • NON-DROWSY
Cold & Flu

NIGHTTIME
Cold & Flu

DAY & NIGHT PACK

NDC 0383-1732-02



Compare to the active ingredients in
Vicks® DayQuil® & Vicks® NyQuil®^{1H}



**DAYTIME
NON-DROWSY
Cold & Flu**

ACETAMINOPHEN / ACHES /
FEVER / SORE THROAT
DEXTROMETHORPHAN HBr /
COUGH
PHENYLEPHRINE HCl /
NASAL CONGESTION

Multi-Symptom

- Pain reliever, fever reducer,
cough suppressant
& nasal decongestant.
- Antihistamine free
- Alcohol free

**NIGHTTIME
Cold & Flu**

ACETAMINOPHEN / ACHES /
FEVER / SORE THROAT
DEXTROMETHORPHAN HBr /
COUGH
DOXYLAMINE SUCCINATE /
SNEEZING / RUNNY NOSE

Multi-Symptom

- Pain reliever, fever reducer,
cough suppressant
& antihistamine
- ALCOHOL 10%

2 – 12 FL OZ (355 mL) BOTTLES / TOTAL 24 FL OZ (1.5 pt) (710 mL)



Cherry
flavor

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION
GUARANTEED

ITEM 984380 WY0106-0722-L



Gluten Free



**PAPER
BOX**



Empty &
Replace Cap
**PLASTIC
BOTTLE**



**PLASTIC
CUP**

how2recycle.info

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

¹Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
²These products are not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks® NyQuil® and DayQuil®.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION
GUARANTEED

ITEM 984380 WY0106-0722-L



DO NOT USE IF PRINTED NECKBAND IS
BROKEN OR MISSING

Gluten Free



PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

¹Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

²These products are not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks® NyQuil® and DayQuil®.

Nighttime Cold & Flu

Drug Facts

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 850 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine

Uses temporarily relieves common cold/flu symptoms:
 ■ cough due to minor throat and bronchial irritation
 ■ sore throat ■ headache ■ minor aches and pains
 ■ fever ■ runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease ■ glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur ■ avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning
 - only use the dose cup provided ■ do not exceed 4 doses per 24 hrs
- | | |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 6 hrs |
| children 4 to under 12 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- each 30 mL contains: sodium 39 mg
- store at 20-25°C (68-77°F)

Inactive ingredients alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments? 1-800-719-9260

Nighttime Cold & Flu

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion ■ cough due to minor throat and bronchial irritation
- sore throat ■ headache ■ minor aches and pains ■ fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen

Adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease ■ heart disease ■ high blood pressure
- thyroid disease ■ diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning
 - only use the dose cup provided ■ do not exceed 4 doses per 24 hrs
- | | |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs | 15 mL every 4 hrs |
| children 4 to under 6 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- each 15 mL contains: sodium 7 mg ■ store at 20-25°C (68-77°F)

Inactive ingredients butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments? 1-800-719-9260




DAY & NIGHT PACK

NDC 0363-0677-02



Compare to the active ingredients in
Vicks® DayQuil® Severe Cold & Flu
& Vicks® NyQuil® Severe Cold & Flu††



**DAYTIME
NON-DROWSY**

Severe Cold & Flu

ACETAMINOPHEN / ACHES /
FEVER / SORE THROAT
DEXTROMETHORPHAN HBr / COUGH
GUAIFENESIN / CHEST CONGESTION
PHENYLEPHRINE HCl /
NASAL CONGESTION

Maximum Strength

- Pain reliever, fever reducer,
cough suppressant, expectorant
& nasal decongestant
- Antihistamine free
- Alcohol free

NIGHTTIME

Severe Cold & Flu

ACETAMINOPHEN / ACHES / FEVER /
SORE THROAT
DEXTROMETHORPHAN HBr / COUGH
DOXYLAMINE SUCCINATE /
SNEEZING / RUNNY NOSE
PHENYLEPHRINE HCl /
NASAL & SINUS CONGESTION /
SINUS PRESSURE

Maximum Strength

- Pain reliever, fever reducer,
cough suppressant,
antihistamine &
nasal decongestant
- ALCOHOL 10%

2 - 12 FL OZ (355 mL) BOTTLES / TOTAL 24 FL OZ (1.5 pt) (710 mL)

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015



**100% SATISFACTION
GUARANTEED**

walgreens.com
©2022 Walgreen Co.

ITEM 908029 W10106-0722-L



3 11917 17820 2

Gluten Free



PAPER
BOX



PLASTIC
BOTTLE



PLASTIC
CUP

how2recycle.info

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

†Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
††These products are not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks® NyQuil® and DayQuil®.

Nighttime Severe Cold & Flu

Drug Facts

Active Ingredients (in each 30 mL)

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Sleeping aid
Phenylephrine HCl 10 mg	Nasal decongestant

Purpose

Uses: temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and sneezing
- cough due to minor throat and bronchial irritation
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores free breathing through the nose

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- hives
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use:

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have:

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent chronic cough as occurs with smoking, asthma, or emphysema
- a doctor-restricted diet

Ask a doctor or pharmacist before use if you are:

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product:

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if:

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Do not take any more if you do not notice any signs or symptoms.

Directions

Take only as directed - see Overdose warning.

Only use the dose cup provided. Do not exceed 4 doses per 24 hrs.

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 30 mL contains sodium 44 mg
- store at 20-25°C (68-77°F)

Inactive ingredients: alcohol, anhydrous citric acid, D&G yellow #10, edetate disodium, FD&C green #3, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

Questions or comments? 1-800-719-8280

Nighttime Severe Cold & Flu

Drug Facts

Active Ingredients (in each 15 mL)

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guafenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Purpose

Uses: temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores free breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to aid the bronchial passageways of both mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- takes with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- hives
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use:

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have:

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not use more than directed.

Stop use and ask a doctor if:

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Take only as directed - see Overdose warning.

Only use the dose cup provided. Do not exceed 4 doses per 24 hrs.

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains sodium 6 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients: butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthyl, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments? 1-800-719-8280

W3ORG0422-F

NDC 0363-5001-40

Walgreens

Compare to the active ingredients in
Vicks® DayQuil® Severe Cold & Flu™

DAYTIME • NON-DROWSY

Severe Cold & Flu

**ACETAMINOPHEN / PAIN RELIEVER /
FEVER REDUCER
PHENYLEPHRINE HCl /
NASAL DECONGESTANT
DEXTROMETHORPHAN HBr /
COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT**

Maximum Strength

- Headache, fever, sore throat, minor aches & pains
- Chest congestion, thins & loosens mucus
- Nasal/sinus congestion, sinus pressure
- Cough


Honey flavor

12 FL OZ (355 mL)

: 53U40 94 F1

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2022 Walgreen Co.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

ITEM 714456
W10107-0623-F
REV-0323
DO NOT USE IF PRINTED BACKGROUND IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses: temporarily relieves common cold/flu symptoms: ■ sinus congestion and pressure ■ nasal congestion ■ minor aches and pains ■ headache ■ sore throat ■ fever ■ cough due to minor throat and bronchial irritation ■ reduces swelling of nasal passages ■ promotes nasal and/or sinus drainage ■ temporarily restores freer breathing through the nose ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away

PEEL BACK AT CORNER FOR MORE INFORMATION

: 53U40 94 B2

Drug Facts
(continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ if you have ever had an allergic reaction to this product or any of its ingredients

HESIVE AREA • NO VARNISH • NO TYPE

Drug Facts
(continued)

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed - see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor

ADHESIVE AREA • NO VARNISH • NO TYPE

AD
NO COATING

Ask a doctor before use if you have

- liver disease ■ heart disease
- high blood pressure ■ diabetes
- thyroid disease ■ cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

◀

3164

children under 4 yrs do not use

Other information ■ each 15 mL contains: sodium 15 mg ■ store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C red #40, FD&C yellow #6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions? 1-800-719-9260

¹Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

²This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks[®] and DayQuil[®].

NO COATING



DO NOT PRINT
Lot and Exp ONLY

Drug Facts (continued)
magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, propylene glycol, stearic acid, silicon dioxide, stearic acid, titanium dioxide
(Questions or comments? 1-800-426-9391)

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Drug Facts (continued)
prostate gland ■ glaucoma ■ thyroid disease
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers
When using this product
do not exceed recommended dosage
excitability may occur, especially in children
alcohol, sedatives, and tranquilizers may increase drowsiness
use caution when driving a motor vehicle or operating machinery ■ avoid alcoholic beverages
Stop use and ask a doctor if
pain or nasal congestion gets worse or lasts more than 7 days ■ new symptoms occur
fever gets worse or lasts more than 3 days
redness or swelling is present
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions ■ do not take more than directed
adults and children 12 years and over
take 2 gelcaps every 4 hours
do not take more than 10 gelcaps in 24 hours
children under 12 years: ask a doctor
Other information
contains FD&C Yellow #5 (tartrazine) as a color additive
Inactive ingredients corn starch, croscarmellose sodium, crospovidone, FD&C red #3, FD&C red #40, FD&C yellow #5, FD&C yellow #6, gelatin, hydroxypropyl cellulose, hydroxypropyl methylcellulose, iron oxide black, iron oxide red, iron oxide yellow, ■ see end flap for expiration date and lot number

Drug Facts (continued)
tamponade relieves these additional symptoms of hay fever: ■ itchy, watery eyes
itching of the nose or throat
helps clear nasal passages
helps decongest sinus openings and passages
Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen
3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
skin redness ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product
If you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if you have
a breathing problem such as emphysema or chronic bronchitis ■ high blood pressure ■ heart disease ■ diabetes ■ liver disease ■ difficulty in urination due to enlargement of the prostate

Drug Facts (continued)
TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
BROKEN ■ see end flap for expiration date and lot number
Drug Facts (continued)
avoid high humidity
TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
BROKEN ■ see end flap for expiration date and lot number

Our pharmacists recommend the Walgreens brand.
how2recycle.info
PAPER BOX
MULTI-LAYER TRAY
DISTRIBUTED BY:
WALGREEN CO.
200 WILMOT RD.,
DEERFIELD, IL 60015
100% SATISFACTION
GUARANTEED
walgreens.com
©2021 Walgreen Co.

ITEM 510071 W00000-0000-0
3 11917 10374 7
50844
REV0422A55908



No print area
Lot no/Exp date

B-2201-453-07-HR
ORG082045307

NDC 0363-4531-07

Walgreens

Compare to Vicks[®] NyQuil[™]
Severe Nighttime Cold &
Flu Relief active ingredients^{††}

NDC 0363-0331-26

NIGHTTIME

Severe Cold & Flu

ACETAMINOPHEN / FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
DOXYLAMINE SUCCINATE / ANTIHISTAMINE
PHENYLEPHRINE HCl / NASAL DECONGESTANT

MAXIMUM STRENGTH

- Pain reliever, fever reducer,
cough suppressant, antihistamine
& nasal decongestant

ORIGINAL
FLAVOR

8 FL OZ (237 mL)

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN

Drug Facts

Active ingredients

(in each 30 mL dose cup or 2 tablespoons)

Acetaminophen 650 mg.....Pain reliever/Fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Doxylamine Succinate 12.5 mg.....Antihistamine
Phenylephrine HCl 10 mg.....Nasal decongestant

Purpose

Uses Temporarily relieves cold/flu symptoms: ■ nasal congestion ■ sinus congestion & pressure ■ sore throat ■ headache ■ minor aches and pain ■ runny nose and sneezing ■ temporarily reduces fever ■ cough due to minor throat and bronchial irritation ■ promotes nasal and/or sinus drainage

Warnings **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use: ■ with any other drug containing

Drug Facts (cont'd)

acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ to make a child sleep.

Ask a doctor before use if you have ■ a sodium restricted diet ■ liver disease ■ heart disease ■ glaucoma ■ high blood pressure ■ thyroid disease ■ diabetes ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema ■ trouble urinating due to enlarged prostate gland.

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED walgreens.com ©2020 Walgreen Co.

LR-151

ITEM 268947 W10139-1120-F



ORG1020-F

Drug Facts (cont'd)

Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin.

When using this product ■ do not use more than directed ■ avoid alcoholic drinks ■ marked drowsiness may occur ■ excitability may occur, especially in children; ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ redness or swelling is present ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as recommended. See Warnings
- use dose cup or teaspoon (TBS)
- do not exceed 4 doses per 24 hours

Age	Dose
Adults & children 12 years and over	30 mL (2 TBS) every 4 hours
Children 4 to under 12 years	Ask a doctor
Children under 4 years	Do not use

Other information

- each 30 mL contains: sodium 45 mg
- store at room temperature

Inactive ingredients anhydrous citric acid, D&C Yellow # 10, FD&C Green # 3, FD&C Yellow # 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sorbitol, xanthan gum

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

Walgreens Pharmacy is a registered trademark of Walgreen Company. ©2020 Walgreen Company. This product is not manufactured or distributed by Walgreen Pharmacy. See back of bottle for more information.

NDC 0363-2029-40

Walgreens

Compare to the active ingredients in
Vicks® NyQuil® Severe Cold & Flu™

**NIGHTTIME
Severe
Cold & Flu**

**ACETAMINOPHEN / PAIN RELIEVER /
FEVER REDUCER
PHENYLEPHRINE HCl /
NASAL DECONGESTANT
DEXTROMETHORPHAN HBr /
COUGH SUPPRESSANT
DOXYLAMINE SUCCINATE /
ANTIHISTAMINE**

Maximum Strength

- Headache, fever, sore throat, minor aches & pains
- Nasal/sinus congestion, sinus pressure
- Sneezing & runny nose
- Cough • Cold & flu


Honey flavor

12 FL OZ (355 mL)

: 6LS40 94 F1

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2022 Walgreen Co.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

ITEM 714663
W10107-0623-F
REV-0323
DO NOT USE IF POINTED NECKBAND IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Doxylamine Succinate 6.25 mg.....	Antihistamine
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses: temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion and pressure ■ cough due to minor throat and bronchial irritation ■ cough to help you sleep ■ minor aches and pains ■ headache ■ sore throat ■ fever ■ runny nose and sneezing ■ reduces swelling of nasal passages ■ temporarily restores freer breathing through the nose ■ promotes nasal and/or sinus drainage

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

PEEL BACK AT CORNER FOR MORE INFORMATION

: 6LS40 94 B2

Drug Facts (continued)

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients ■ to make a child sleepy

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ glaucoma ■ thyroid disease ■ diabetes ■ cough that occurs with too much

ADHESIVE AREA
NO VARNISH • NO TYPE

Drug Facts (continued)

■ avoid alcoholic drinks ■ be careful when driving a motorvehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed - see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

ADHESIVE AREA
NO VARNISH • NO TYPE

ADH
NO COATING

pregn (mucus) ■ trouble urinating due to an enlarged prostate gland
 ■ a breathing problem such as emphysema or chronic bronchitis
 ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

When using this product
 ■ do not use more than directed
 ■ excitability may occur, especially in children
 ■ marked drowsiness may occur

◀

NO COATING



adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information ■ each 15 mL contains: sodium 15 mg ■ store at 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C red #40, FD&C yellow #6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions?
 1-800-719-9260

¹⁰Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
¹¹This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks® and NyQuil®.



NDC 0363-0603-40

Walgreens

Compare to the active ingredients in
Vicks® DayQuil® Severe Cold & Flu[†]

DAYTIME • NON-DROWSY

Severe Cold & Flu

ACETAMINOPHEN /
ACHES / FEVER / SORE THROAT
DEXTROMETHORPHAN HBr / COUGH
GUAIFENESIN / CHEST CONGESTION
PHENYLEPHRINE HCl /
NASAL CONGESTION

Maximum Strength

- Pain reliever, fever reducer,
cough suppressant, expectorant
& nasal decongestant
- Antihistamine free
- Alcohol free

12 FL OZ (355 mL)

: 60340 94 F7

DISTRIBUTED BY: WALGREEN CO.
300 WALSH RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com 43821 Walgreen Co.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

ITEM 330479
W10107-0622-F
Gluten Free
W30RG0222-F

DO NOT USE IF PRINTED NECKBAND IS CRUMPLED OR MISSING

Drug Facts

Active Ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/ fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses: temporarily relieves common cold/flu symptoms: ■ sinus congestion and pressure ■ nasal congestion ■ minor aches and pains ■ headache ■ sore throat ■ fever ■ cough due to minor throat and bronchial irritation ■ reduces swelling of nasal passages ■ promotes nasal and/or sinus drainage ■ temporarily restores freer breathing through the nose ■ helps loosen phlegm (mucus) and thin bronchial secretions to aid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

PEEL BACK AT CORNER FOR MORE INFORMATION

1191716340

: 60340 94 B1

Drug Facts
(continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ cough that occurs with too much phlegm (mucus)

ADHESIVE AREA
NO COATING • NO VARNISH • NO TYPE

Drug Facts
(continued)

When using this product do not use more than directed

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed — see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information ■ each 15 mL contains:

ADHESIVE AREA
NO COATING • NO VARNISH • NO TYPE